

Actions of compact groups on coherent sheaves

Jürgen Hausen and Peter Heinzner¹

Let K be a compact Lie group with complexification $K^{\mathbb{C}}$ and X a reduced Stein space endowed with a continuous action of K by holomorphic transformations. In this set up there is a complex space $X//K$ and a surjective K -invariant holomorphic map $\pi: X \rightarrow X//K$ such that the structure sheaf of $X//K$ is the sheaf of invariants $(\pi_*\mathcal{O}_X)^K$. Furthermore, the K -action on X can be complexified in the following sense: X can be realized as an open K -stable subset of a Stein space X^c which is endowed with a holomorphic $K^{\mathbb{C}}$ -action. Moreover, the inclusion $X \subset X^c$ is universal, i.e., every K -equivariant holomorphic map ϕ from X into a holomorphic $K^{\mathbb{C}}$ -space Z extends uniquely and $K^{\mathbb{C}}$ -equivariantly to a holomorphic map $\phi^c: X^c \rightarrow Z$ (see [H]).

In the present paper we extend the above results to Stein spaces X which are not necessarily reduced. More importantly, we show that every continuous coherent K -sheaf over X can be extended to a coherent $K^{\mathbb{C}}$ -sheaf over X^c . This is also proved in the context of non-reduced spaces.

The constructions of quotients and complexifications are compatible with the reduced structures. Thus we have $\text{red}(X)//K = \text{red}(X//K)$ and $\text{red}(X)^c = \text{red}(X^c)$, where red associates to a complex space its underlying reduced space. Moreover, the results are also valid for K -spaces X with $\text{red}(X)$ admitting a semistable quotient (see Section 1). In this context we show that the sheaf $(\pi_*\mathcal{S})^K$ of invariant sections of a coherent K -sheaf $\mathcal{S}$ over X is a coherent analytic sheaf over the quotient space $X//K$ (see Section 1). Two direct applications of our results are

- a) For every closed complex K -subspace A of X the image $\pi(A)$ is a closed subspace of $X//K$ such that $A//K \cong \pi(A)$. Moreover, A extends to a closed $K^{\mathbb{C}}$ -subspace A^c of X^c .
- b) Every holomorphic K -vector bundle over X is given by the restriction of a holomorphic $K^{\mathbb{C}}$ -vector bundle over the complexification X^c .

The authors would like to thank the referees for their interest in the subject and for valuable suggestions.

1. Formulation of the results

In this paper a complex space X is a not necessarily reduced complex space with countable topology. By $\text{red}(X)$ we denote the underlying reduced space, i.e., set-theoretically $\text{red}(X) = X$ and $\mathcal{O}_{\text{red}(X)} = \mathcal{O}_X/\mathcal{N}_X$ on the level of sheaves. Here $\mathcal{N}_X$ denotes the nilradical of the structure sheaf $\mathcal{O}_X$ of X .

¹ Supported by a Heisenberg Stipendium of the Deutsche Forschungsgemeinschaft

Let G be a real Lie group and assume that G acts continuously on X by holomorphic transformations. Then G acts also on the structure sheaf $\mathcal{O} := \mathcal{O}_X$ of X : A given $g \in G$ induces for every $x \in X$ an isomorphism $\mathcal{O}_x \rightarrow \mathcal{O}_{g \cdot x}$, $f \mapsto g \cdot f$, of stalks. We call X a *complex G -space* if for all open sets $N \subset G$ and $U_1, U_2 \subset X$ with $U_2 \subset \bigcap_{g \in N} g \cdot U_1$ the map

$$(1) \quad \Phi: N \times \mathcal{O}(U_1) \rightarrow \mathcal{O}(U_2), \quad (g, f) \mapsto (g \cdot f)|_{U_2}$$

is continuous with respect to the canonical Frechét topology on $\mathcal{O}(U_j)$, $j = 1, 2$ (see [G-R], Chapter V, §6). If G is a complex Lie group and all above maps Φ are holomorphic, then we say that X is a *holomorphic G -space*.

Now let $\mathcal{S}$ be a G -sheaf on X , i.e., G acts on $\mathcal{S}$ such that the projection $\mathcal{S} \rightarrow X$, $s_x \mapsto x$ is G -equivariant. Reformulating (1) for a coherent sheaf $\mathcal{S}$, we obtain the notion of a *continuous* (resp. *holomorphic*) *coherent G -sheaf* on X .

A holomorphic map $\phi: X \rightarrow X'$ of two complex G -spaces X and X' is called *G -equivariant*, if it is G -equivariant as a map of sets and the comorphism ϕ^0 is compatible with the actions of G on the structure sheaves $\mathcal{O}_X$ and $\mathcal{O}_{X'}$. The map ϕ is called *G -invariant* if it is equivariant with respect to the trivial action of G on X' .

Remark 1. If X is reduced, then it is a complex G -space. If moreover G is a complex Lie group and the action $G \times X \rightarrow X$ is holomorphic, then X is a holomorphic G -space (see [K]). A holomorphic map of reduced G -spaces is equivariant if and only if it is equivariant as a map of sets.

Now let K be a compact real Lie group and assume that X is a complex K -space. A K -invariant holomorphic map $\pi: X \rightarrow X//K$ onto a complex space $X//K$ is said to be a *semistable quotient* of the K -space X , if

- i) the structure sheaf $\mathcal{O}_{X//K}$ of $X//K$ is the sheaf of invariants $(\pi_* \mathcal{O}_X)^K$, i.e., for every open set $Q \subset X//K$ we have $\mathcal{O}_{X//K}(Q) = \mathcal{O}_X(\pi^{-1}(Q))^K$ and
- ii) π is a Stein map, i.e., for every Stein open set $Q \subset X//K$ the inverse image $\pi^{-1}(Q)$ is also Stein.

The above definition generalizes the notion introduced in [H-M-P] to the non-reduced case. Examples of semistable quotients occur in Geometric Invariant Theory. Our first main result is the following

Quotient Theorem. *A semistable quotient $\pi: X \rightarrow X//K$ for X exists if and only if it exists for $\text{red}(X)$. If this is the case, then $\text{red}(\pi): \text{red}(X) \rightarrow \text{red}(X//K)$ is a semistable quotient for $\text{red}(X)$.*

By [H], Section 6.5, every reduced Stein K -space has a semistable quotient and the associated quotient space is again Stein. This, together with the above result, implies the following

Corollary 1. *If X is a Stein K -space, then the semistable quotient $\pi: X \rightarrow X//K$ exists. Moreover, $X//K$ is a Stein space.* $\square$

Since the reduction of a semistable quotient is a semistable quotient of the associated reduced K -space, [H], 2.3, yields that semistable quotients are universal with respect to K -invariant holomorphic maps. In particular, the quotient space $X//K$ is unique up to isomorphism.

If X admits a semistable quotient $\pi: X \rightarrow X//K$, then we denote for a K -sheaf $\mathcal{S}$ on X by $(\pi_*\mathcal{S})^K$ the sheaf of invariants on $X//K$, i.e., $(\pi_*\mathcal{S})^K(Q) := \mathcal{S}(\pi^{-1}(Q))^K$. The following fact is a generalization of a result of Roberts (see [R]) for holomorphic $K^\mathbb{C}$ -sheaves.

Coherence Theorem. *Assume that X has a semistable quotient $\pi: X \rightarrow X//K$ and let $\mathcal{S}$ be a continuous coherent K -sheaf on X . Then the sheaf $(\pi_*\mathcal{S})^K$ on $X//K$ is coherent.*

Let $K^\mathbb{C}$ denote the complexification of K . A *complexification* X^c of a complex K -space X is a holomorphic $K^\mathbb{C}$ -space X^c which contains X as a K -stable open subset such that every K -equivariant holomorphic map ϕ from X into a holomorphic $K^\mathbb{C}$ -space Z extends uniquely to a $K^\mathbb{C}$ -equivariant holomorphic map $\phi^c: X^c \rightarrow Z$. Note that a complexification X^c is unique up to isomorphism and that $X^c = K^\mathbb{C} \cdot X$.

Complexification Theorem. *If X has a semistable quotient $\pi: X \rightarrow X//K$, then the complexification X^c exists. Moreover, the extension $\pi^c: X^c \rightarrow X//K$ is a semistable quotient for the $K^\mathbb{C}$ -space X^c and $\text{red}(X^c) = \text{red}(X)^c$ holds.*

This result generalizes the Theorem in Section 6.6 of [H], where the existence of complexifications is proved for reduced Stein K -spaces.

Now, let $\mathcal{S}$ be a continuous coherent K -sheaf on X and assume that X has a semistable quotient $\pi: X \rightarrow X//K$. A holomorphic $K^\mathbb{C}$ -sheaf $\mathcal{S}^c$ on X^c with $\mathcal{S}^c|_X = \mathcal{S}$ is called a $K^\mathbb{C}$ -extension of $\mathcal{S}$.

Extension Theorem. *Assume that X has a semistable quotient and let $\mathcal{S}$ be a continuous coherent K -sheaf $\mathcal{S}$ on X . Then, up to $K^\mathbb{C}$ -equivariant isomorphism, there is a unique $K^\mathbb{C}$ -extension of $\mathcal{S}$.*

For a K -stable closed subspace A of X which is defined by a K -invariant sheaf $\mathcal{I}$ of ideals, the Coherence and Extension Theorem imply:

Corollary 2. *The sheaf $(\pi_*\mathcal{I})^K$ of ideals endows $\pi(A)$ with the structure of closed subspace of $X//K$. The restriction of π to A is a semistable quotient for A . Moreover, A has a complexification A^c by a closed $K^\mathbb{C}$ -subspace of X^c and $\pi(A) = \pi^c(A^c)$. $\square$*

If $\mathcal{S}$ is a locally free continuous K -sheaf over X with $K^\mathbb{C}$ -extension $\mathcal{S}^c$, then the complement E of the set of points in X^c at which $\mathcal{S}^c$ is not locally free is a proper analytic $K^\mathbb{C}$ -stable subset of X^c . Since $X^c = K^\mathbb{C} \cdot X$, this implies that E is empty. In other words we have the following

Corollary 3. *The extension of a locally free K -sheaf over X is a locally free $K^\mathbb{C}$ -sheaf over X^c . In particular, every holomorphic K -vector bundle over X extends to a holomorphic $K^\mathbb{C}$ -vector bundle over X^c . $\square$*

2. Equivariant Resolution

Let K be a compact real Lie group and let X be a complex K -space. In this section we assume that the associated reduced K -space $\text{red}(X)$ has a semistable quotient $\pi_r: \text{red}(X) \rightarrow \text{red}(X)//K$. Let $\mathcal{S}$ be a continuous coherent K -sheaf on X . Note that we have a continuous representation of K on the Frechét space $\mathcal{S}(X)$.

A section $s \in \mathcal{S}(X)$ of $\mathcal{S}$ is said to be K -finite if the vector subspace of $\mathcal{S}(X)$ generated by $K \cdot s$ is of finite dimension. We denote the vector subspace of K -finite elements of $\mathcal{S}(X)$ by $\mathcal{S}(X)_{\text{fin}}$. By a theorem of Harish-Chandra (see [HC], Lemma 5), $\mathcal{S}(X)_{\text{fin}}$ is dense in $\mathcal{S}(X)$.

Equivariant Resolution Lemma. *Every $y \in \text{red}(X)//K$ has an open neighborhood Q such that over $U := \pi_r^{-1}(Q)$ there is an exact sequence $\mathcal{V} \rightarrow \mathcal{S}|_U \rightarrow 0$ of K -sheaves. Here V is a finite dimensional representation space of K and $\mathcal{V} := \mathcal{O} \otimes V$ is endowed with the action defined by $k \cdot (f \otimes v) := k \cdot f \otimes k \cdot v$.*

Proof. Since π_r is a Stein map, we may assume that X is a Stein space. By Proposition 3.2 and Corollary 2 in Section 2.3 of [H], there is a unique K -invariant analytic set A of minimal dimension contained in $\pi_r^{-1}(y)$. We fix a point $x \in A$. Then, since X is a Stein space, there are sections $s_1, \dots, s_r \in \mathcal{S}(X)$ which generate $\mathcal{S}_x$ as an $\mathcal{O}_x$ -module.

The K -finite elements of $\mathcal{S}(X)$ are dense. Thus we may assume that the s_i generate a finite dimensional K -submodule V of $\mathcal{S}(X)$. Hence there is an equivariant homomorphism $\alpha: \mathcal{V} \rightarrow \mathcal{S}$ of K -sheaves which is defined by $\mathbf{f} \otimes s \rightarrow \mathbf{f} \mathbf{s}$ where $\mathbf{s}$ denotes the section of $\mathcal{S}$ defined by $s \in V \subset \mathcal{S}(X)$.

Since α is a homomorphism of coherent K -sheaves, $B := \text{Supp}(\mathcal{S}/\alpha(\mathcal{V}))$ is a closed K -invariant analytic subset of X . By definition, α is surjective over $X \setminus B$. Furthermore, we have $\pi_r(A) \cap \pi_r(B) = \emptyset$. Hence $Q := \text{red}(X)//K \setminus \pi_r(B)$ is open, contains y and α is surjective over $U := \pi_r^{-1}(Q)$. $\square$

As a consequence of the Equivariant Resolution Lemma we give here a proof of the Coherence Theorem in the reduced Stein case (see also [R]). We use two well known Lemmas. The proof of the first one can be found for example in [R] or [Sch].

Lemma 1. *Let V and W be finite dimensional complex $K^{\mathbb{C}}$ -modules. Then there exist K -equivariant polynomials $p_i: W \rightarrow V$, $i = 1, \dots, r$ which generate $(\pi_{W*} \mathcal{V})^K$ as an $\mathcal{O}_{W//K^{\mathbb{C}}}$ -module.* $\square$

Lemma 2. *For every exact sequence $\mathcal{S}_1 \rightarrow \mathcal{S}_2 \rightarrow \mathcal{S}_3$ of continuous coherent K -sheaves on X the induced sequence $(\pi_{r*} \mathcal{S}_1)^K \rightarrow (\pi_{r*} \mathcal{S}_2)^K \rightarrow (\pi_{r*} \mathcal{S}_3)^K$ on $\text{red}(X)//K$ is also exact.*

Proof. Integration over K defines a projection operator $\mathcal{S}(X) \rightarrow \mathcal{S}(X)^K$. Since exactness is a local property and $\pi_r: \text{red}(X) \rightarrow \text{red}(X)//K$ is a Stein map, the assertion follows. $\square$

Proposition 1. *Let X be a reduced Stein K -space and denote by $\pi: X \rightarrow X//K$ the semistable quotient. Then $(\pi_* \mathcal{S})^K$ is a coherent sheaf on $X//K$.*

Proof. First consider the case $\mathcal{S} = \mathcal{V} := \mathcal{O}_X \otimes V$ with some K -module V . By [H], 6.6, we may assume that X is a holomorphic $K^\mathbb{C}$ -space. Moreover, by [H], 5.4 and 6.2, we may assume that X is a closed K -subspace of an open Stein π_W -saturated K -subspace of some finite dimensional K -module W . So, for the case $\mathcal{S} = \mathcal{V}$, Lemma 1 provides $p_1, \dots, p_r \in \mathcal{S}(X)^K$ that generate $(\pi_*\mathcal{S})^K$, i.e., the associated sequence

$$\mathcal{O}_{X//K}^r \xrightarrow{\alpha} (\pi_*\mathcal{S})^K \rightarrow 0.$$

is exact. The kernel of α is $(\pi_*\mathcal{R})^K$, where $\mathcal{R} \subset \mathcal{O}_X^r$ denotes the K -sheaf of relations of the generators p_i . Applying Lemma 2 to an equivariant resolution of $\mathcal{R}$ yields that also $\text{Ker}(\alpha)$ is finitely generated. Consequently $(\pi_*\mathcal{S})^K$ is coherent. For general $\mathcal{S}$, apply Lemma 2 to an equivariant resolution $\mathcal{V}' \rightarrow \mathcal{V} \rightarrow \mathcal{S} \rightarrow 0$. $\square$

3. $K^\mathbb{C}$ -Extensions

In this section we apply the Equivariant Resolution Lemma to investigate extensions of coherent K -sheaves. Assume that X is a complex K -space such that there is a semistable quotient $\pi_r: \text{red}(X) \rightarrow \text{red}(X)//K$ and a complexification X^c with the following properties:

- i) $\text{red}(X^c)$ is the complexification of $\text{red}(X)$ and $\pi_r^c: \text{red}(X^c) \rightarrow \text{red}(X)//K$ is a semistable quotient.
- ii) For every open subspace of the form $U := \pi_r^{-1}(Q)$ with $Q \subset \text{red}(X)//K$ open, the open subspace $U^c := K^\mathbb{C} \cdot U = (\pi_r^c)^{-1}(Q)$ of X^c is a complexification of U .

Note that, by [H], Sections 3.3 and 6.6, these assumptions are valid if X is a reduced Stein K -space.

Identity Principle. *Let $\mathcal{T}$ be a holomorphic coherent $K^\mathbb{C}$ -sheaf over X^c . Then the restriction map $R: \mathcal{T}(X^c)_{\text{fin}} \rightarrow \mathcal{T}(X)_{\text{fin}}$ is bijective.*

Proof. First we show that R is injective. So, let $s \in \mathcal{T}(X^\mathbb{C})$ be K -finite such that $s|_X = 0$. Since $u|_X = 0$ for all $u \in V := \text{Lin}_\mathbb{C}(K \cdot s) = \text{Lin}_\mathbb{C}(K^\mathbb{C} \cdot s)$, we have $(g \cdot s)|_X = 0$ for all $g \in K^\mathbb{C}$. Hence $s|_{g^{-1} \cdot X} = 0$ for every $g \in K^\mathbb{C}$. Since $X^c = K^\mathbb{C} \cdot X$, it follows that $s = 0$.

In order to show that R is surjective, we first assume that X is Stein and that there is an equivariant resolution $\mathcal{V} \xrightarrow{\alpha} \mathcal{T} \rightarrow 0$ with V a finite-dimensional K -module and $\mathcal{V} := \mathcal{O}_{X^c} \otimes V$. Every K -finite $F \in \mathcal{V}(X)$ defines a finite-dimensional K -module $W := \text{Lin}_\mathbb{C}(K \cdot f)$.

Evaluating the elements of W determines a K -equivariant holomorphic map $\Phi: X \rightarrow \text{Hom}(W, V)$. Since Φ extends to a $K^\mathbb{C}$ -equivariant holomorphic map $\Phi^c: X^c \rightarrow V$, it follows that F extends to X^c as a holomorphic map. This yields surjectivity of the restriction $\mathcal{V}(X^c)_{\text{fin}} \rightarrow \mathcal{V}(X)_{\text{fin}}$.

Surjectivity of R is obtained as follows. Let $s \in \mathcal{T}(X)_{\text{fin}}$. Since X was assumed to be Stein, we find an $s_1 \in \mathcal{V}(X)$ with $\alpha(s_1) = s$. Note that α maps $\text{Lin}_\mathbb{C}(K \cdot s_1)$ onto the finite-dimensional vector space $\text{Lin}_\mathbb{C}(K \cdot s)$. Hence we may assume that s_1 is K -finite. As seen above, s has an extension $s_1^c \in \mathcal{V}(X^c)_{\text{fin}}$. Then $\alpha(s_1^c) \in \mathcal{T}(X^c)_{\text{fin}}$ is an extension of s .

Now, in the general case, let $s \in \mathcal{T}(X)_{\text{fin}}$. The Equivariant Resolution Lemma and the above consideration yield a cover of $\text{red}(X//K)$ by open sets Q_i such that we can extend s over each $U_i = \pi_r^{-1}(Q_i)$ to $s_i^c \in \mathcal{T}(U_i^c)_{\text{fin}}$. By injectivity of R , we obtain that any two such extensions s_i^c and s_j^c coincide over $U_i^c \cap U_j^c$. Thus the s_i^c patch together to an extension $s^c \in \mathcal{T}(X^c)$ of s . As before we can achieve that s^c is K -finite. $\square$

Let $\mathcal{S}^1$ and $\mathcal{S}^2$ be holomorphic coherent $K^{\mathbb{C}}$ -sheaves on X^c . Then $\mathfrak{H}om(\mathcal{S}^1, \mathcal{S}^2)$ is a holomorphic $K^{\mathbb{C}}$ -sheaf. The action of $g \in K^{\mathbb{C}}$ on $\mathbf{F} \in \mathfrak{H}om(\mathcal{S}^1, \mathcal{S}^2)_x$ is given by

$$(g \cdot \mathbf{F})(\mathbf{s}) := g \cdot (\mathbf{F}(g^{-1} \cdot \mathbf{s})).$$

where $\mathbf{s} \in \mathcal{S}_{g \cdot x}^1$. The $K^{\mathbb{C}}$ -invariant global sections of $\mathfrak{H}om(\mathcal{S}^1, \mathcal{S}^2)$ are precisely the $K^{\mathbb{C}}$ -equivariant homomorphisms from $\mathcal{S}^1$ to $\mathcal{S}^2$. So the Identity Principle yields:

Homomorphism Lemma. *Every homomorphism $\alpha : \mathcal{S}^1|_X \rightarrow \mathcal{S}^2|_X$ of K -sheaves extends uniquely to a homomorphism $\alpha^c : \mathcal{S}^1 \rightarrow \mathcal{S}^2$ of $K^{\mathbb{C}}$ -sheaves.* $\square$

Local Extension Lemma. *Let $\mathcal{S}$ be a continuous coherent K -sheaf on X . Then every $y \in \text{red}(X)//K$ has an open neighborhood Q such that the restriction $\mathcal{S}_U$ of $\mathcal{S}$ to $U := \pi_r^{-1}(Q)$ has a $K^{\mathbb{C}}$ -extension $\mathcal{S}_U^c$.*

Proof. By the Equivariant Resolution Lemma we find an open Stein neighborhood $Q \subset \text{red}(X)//K$ of y such that over U there exists an equivariant resolution $\mathcal{V}_1 \xrightarrow{\alpha} \mathcal{V}_2 \rightarrow \mathcal{S}_U \rightarrow 0$, where the V_i are finite-dimensional representation spaces of K and $\mathcal{V}_i$ are the associated K -sheaves on X .

Set $\mathcal{V}_i^c := \mathcal{O}_{X^c} \otimes V_i$ and endow each $\mathcal{V}_i^c$ with the diagonal $K^{\mathbb{C}}$ -action. By the Homomorphism Lemma, there is a unique $K^{\mathbb{C}}$ -equivariant extension $\alpha^c : \mathcal{V}_1^c \rightarrow \mathcal{V}_2^c$ of α . Identify $\mathcal{S}_U$ with $\mathcal{V}_2/\text{Im}(\alpha)$ and set $\mathcal{S}_U^c := \mathcal{V}_2^c/\text{Im}(\alpha^c)$. $\square$

4. Invariant subspaces of reduced spaces

Let K be a compact real Lie group and X a complex K -space. Assume that the associated reduced K -space $\text{red}(X)$ has a semistable quotient $\pi_r : \text{red}(X) \rightarrow \text{red}(X)//K$. A technical ingredient for the proofs of our results is the following

Local Embedding Lemma. *Every point $y \in (\text{red } X)//K$ has an open Stein neighborhood $Q \subset (\text{red } X)//K$ such that the open subspace $U := \pi_r^{-1}(Q)$ of X can be realized as a closed K -subspace of a reduced Stein K -space.*

Proof. We may assume that X is a Stein space. As in the reduced case (see [H], 6.2), we can find a K -equivariant holomorphic map ϕ from X into a complex finite dimensional K -module V such that ϕ is an immersion along the fiber $\pi_r^{-1}(y)$ and $\text{red}(\phi)$ embeds some π_r -saturated open neighborhood $\text{red}(U)$ of x properly into an open K -stable subset Z' of V .

By Siu's Theorem (see [S]), Z' contains a Stein open neighborhood Z'' of $\phi(U)$. Set $Z := \bigcap_{k \in K} k \cdot Z''$. Then Z is an open K -stable Stein neighborhood of $\phi(U)$. Now, the set $A \subset U$ consisting of all points $a \in U$ for which ϕ is not an immersion is a K -stable analytic subset of $\text{red}(U)$ which does not intersect $\pi_r^{-1}(y)$. Hence we may shrink U and Z such that they are still Stein and $\phi|_U: U \rightarrow Z$ is a closed embedding. $\square$

In the sequel, let Z be a reduced Stein K -space and assume that X is a K -subspace of Z . Then X is defined by a continuous coherent K -sheaf $\mathcal{I}$ of ideals in $\mathcal{O}_Z$. Let $\kappa: Z \rightarrow Z//K$ denote the semistable quotient.

Proposition 2. *The ideal $(\kappa_*\mathcal{I})^K$ defines a closed subspace $X//K$ of $Z//K$ and the restriction $\pi := \kappa|_X: X \rightarrow X//K$ is a semistable quotient for X . Moreover, $\text{red}(\pi): \text{red}(X) \rightarrow \text{red}(X//K)$ is a semistable quotient for $\text{red}(X)$.*

Proof. By Proposition 1, $(\kappa_*\mathcal{I})^K$ is coherent and hence $X//K$ is a complex subspace of $Z//K$. The fact that π and $\text{red}(\pi)$ are semistable quotients now follows from applying Lemma 2 to the following two exact sequences of K -sheaves:

$$0 \rightarrow \mathcal{I} \rightarrow \mathcal{O}_Z \rightarrow \mathcal{O}_X \rightarrow 0, \quad 0 \rightarrow \sqrt{\mathcal{I}} \rightarrow \mathcal{O}_Z \rightarrow \mathcal{O}_{\text{red } X} \rightarrow 0. \quad \square$$

Now assume that the ideal $\mathcal{I}$ has an extension to Z^c by a holomorphic $K^\mathbb{C}$ -ideal $\mathcal{I}^c$ of $\mathcal{O}_{Z^c}$. Note that for the $K^\mathbb{C}$ -subspace X^c of Z defined by $\mathcal{I}^c$ we have $X^c = K^\mathbb{C} \cdot X$. In particular, [H], Section 3.3, implies that $\text{red}(X^c)$ is the complexification of $\text{red}(X)$. This statement holds also for the corresponding non-reduced spaces:

Proposition 3. *Let $Q \subset X//K$ be open and $U := \pi^{-1}(Q)$. Then the open subspace $U^c := K^\mathbb{C} \cdot U$ of X^c is the complexification of U .*

Proof. We have to show that U^c is universal with respect to K -equivariant holomorphic maps $\phi: U \rightarrow Y$ into holomorphic $K^\mathbb{C}$ -spaces Y .

As a set U is of the form $U = U^c \cap X$. In particular, U is orbit-convex (see [H], 3.2) and hence $\text{red}(U^c)$ is the complexification of $\text{red}(U)$. So the map ϕ extends to a $K^\mathbb{C}$ -equivariant continuous map $\phi^c: U^c \rightarrow Y$.

Let $\phi^0: \mathcal{O}_Y \rightarrow \phi_*\mathcal{O}_U$ denote the comorphism of ϕ . Then we define the comorphism $(\phi^c)^0: \mathcal{O}_Y \rightarrow \phi^c_*\mathcal{O}_{U^c}$ as follows: For $y \in Y$ and $x \in U$ with $\phi^c(x) = y$ choose a $g_0 \in K^\mathbb{C}$ with $g_0 \cdot x \in U$ and set

$$(\phi^c)^0(\mathbf{f}) := g_0^{-1} \cdot \phi^0(g_0 \cdot \mathbf{f})$$

for every germ $\mathbf{f} \in \mathcal{O}_Y$ at y . We have to show that this definition does not depend on the choice of g_0 . Set

$$N(x) := \{g \in K^\mathbb{C}; g \cdot x \in U\}.$$

Recall that U is an orbit convex subset of U^c and therefore $N(x)$ is connected mod K (see [H], 1.5, 3.2 and 6.6). Now, let $\mathbf{h} := g_0 \cdot \mathbf{f}$ and consider the set

$$M := \{g \in N(x); (gg_0^{-1})^{-1} \cdot \phi^0(gg_0^{-1} \cdot \mathbf{h}) = \phi^0(\mathbf{h})\}.$$

Then $Kg_0 \subset M$ holds. Moreover, M is closed in $N(x)$ because the $K^{\mathbb{C}}$ -sheaves $\mathcal{O}_{X^c}$ and $\mathcal{O}_Y$ are continuous. We claim that M is also open in $N(x)$. Using the holomorphy of the $K^{\mathbb{C}}$ -sheaves $\mathcal{O}_{X^c}$ and $\mathcal{O}_Y$ this is seen as follows: If $g_1 \in M$ and $g_2 := g_1 g_0^{-1}$, then

$$(2) \quad \phi^0(\mathbf{h}) = (gg_2)^{-1} \cdot \phi^0(gg_2 \cdot \mathbf{h})$$

holds for all $g \in K$. After representing $\mathbf{h}$ by some section h defined in a neighborhood of $\phi(g_0 \cdot x)$ and restricting (2) to a sufficiently small neighborhood of $g_0 \cdot x$, the right hand side of (2) depends holomorphically on g .

Now the Identity Theorem yields that (2) is satisfied for g in some neighborhood of $e \in K^{\mathbb{C}}$. This implies that M is open in $N(x)$. Consequently $M = N(x)$ holds, i.e., $(\phi^c)^0$ is well defined. $\square$

5. Proof of the Theorems

Proof of the Quotient Theorem. Assume first that a semistable quotient $\pi: X \rightarrow X//K$ exists. Consider the exact sequence $0 \rightarrow \mathcal{N}_X \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{N}_X \rightarrow 0$ of K -sheaves. By Lemma 2, the associated sequence of sheaves of invariants on $X//K$ is also exact. This implies that $\text{red}(\pi): \text{red}(X) \rightarrow \text{red}(X//K)$ is a semistable quotient for $\text{red}(X)$.

Now assume that there is a semistable quotient $\pi_r: \text{red}(X) \rightarrow \text{red}(X)//K$. On the level of topological spaces set $X//K := \text{red}(X)//K$ and endow $X//K$ with the structure sheaf $\mathcal{O}_{X//K} := \pi_{r*}(\mathcal{O}_X)^K$. Then it follows from Proposition 2 and the Local Embedding Lemma that $X//K$ is a complex space. Moreover, by construction, $\pi := \pi_r$, interpreted as a morphism of the ringed spaces X and $X//K$, is the quotient map. $\square$

Proof of the Coherence Theorem. By the Local Embedding Lemma and Proposition 2, we may assume that X is a K -stable closed subspace of a reduced Stein K -space Z . Noting that the trivial extension of $\mathcal{S}$ to Z is a continuous coherent K -sheaf, we obtain the assertion from Proposition 1. $\square$

Proof of the Complexification Theorem. Let $\pi: X \rightarrow X//K$ be the semistable quotient for X . Choose a cover of $X//K$ by Stein open sets Q_i as in the Local Embedding Lemma. According to the Local Extension Lemma we may assume that every $U_i := \pi^{-1}(Q_i)$ satisfies the assumptions of Proposition 3.

Let U_i^c be the complexification of $U_i := \pi^{-1}(Q_i)$. Note that every complexification $(U_i \cap U_j)^c$ of $U_i \cap U_j$ is contained as an open subspace in both, U_i^c and U_j^c . Consequently the U_i^c can be glued together over $X//K$ to a complexification X^c of X .

By construction, $\text{red}(X^c)$ is the complexification of $\text{red}(X)$. So we obtain $\text{red}(\pi^c) = (\text{red } \pi)^c$ for the K -invariant holomorphic $\pi^c: X^c \rightarrow X//K$ extension of π . In particular, π^c is a Stein map and hence a semistable quotient. $\square$

Note that, by construction, the complexification X^c satisfies the technical assumptions i) and ii), made in Section 3.

Proof of the Extension Theorem. By the Homomorphism Lemma, we only have to prove the existence of a $K^{\mathbb{C}}$ -extension. According to the Local Extension Lemma we can cover $X//K$ by open sets Q_i such that on each $U_i := \pi^{-1}(Q_i)$ the restriction $\mathcal{S}_i := \mathcal{S}|_{U_i}$ has an $K^{\mathbb{C}}$ -extension $\mathcal{S}_i^c$ to $U_i^c = (\pi^c)^{-1}(Q_i)$.

The Homomorphism Lemma yields glueing homomorphisms $\mathcal{S}_i^c|_{U_i^c \cap U_j^c} \rightarrow \mathcal{S}_j^c|_{U_i^c \cap U_j^c}$ that extend the identity map $\mathcal{S}_i|_{U_i \cap U_j} \rightarrow \mathcal{S}_j|_{U_i \cap U_j}$. Hence the $\mathcal{S}_i^c$ can be glued together to a $K^{\mathbb{C}}$ -sheaf $\mathcal{S}^c$ on X^c . It is straightforward to check that $\mathcal{S}^c$ is a $K^{\mathbb{C}}$ -extension of $\mathcal{S}$. $\square$

References

- [G-R] Grauert, H.; Remmert, R.: *Theorie der Steinschen Räume*. Heidelberg: Springer 1977
- [HC] Harish-Chandra: *Discrete Series for semisimple Lie groups II*. Acta. Math. 116, 1-111 (1966)
- [H] Heinzner, P.: *Geometric Invariant Theory on Stein Spaces*. Math. Ann. 289, 631-662 (1991)
- [H-M-P] Heinzner, P.; Migliorini, L.; Polito, M.: *Semistable Quotients*. To appear, Annali della Scuola Normale Superiore di Pisa (1997)
- [K] Kaup, W.: *Infinitesimale Transformationsgruppen auf komplexen Räumen*. Math. Ann. 160, 72-92 (1965)
- [R] Roberts, M.: *A Note on Coherent G -Sheaves*. Math. Ann. 275, 573-582 (1986)
- [Sch] Schwarz, G.: *Lifting smooth homotopies of orbit spaces*. Publ. Math. IHES 51, 37-135 (1980)
- [S] Siu, Y.-T.: *Every Stein subvariety admits a Stein neighborhood*. Inv. Math. 38, 89-100 (1976)

Jürgen Hausen
Fakultät für Mathematik und
Informatik, Universität Konstanz
78457 Konstanz
Germany
e-mail: Juergen.Hausen@uni-konstanz.de

Peter Heinzner
Department of Mathematics
Brandeis University
Waltham, MA 02254-9110
USA
e-mail: heinzner@max.math.brandeis.edu